

CellCept[®]

Mycophenolate mofetil (MMF)

1. DESCRIPTION

1.1 THERAPEUTIC / PHARMACOLOGIC CLASS OF DRUG

Selective Immunosuppressive agent – Mycophenolic acid (ATC Code – L04AA06)

1.2 TYPE OF DOSAGE FORM

Oral administration: CellCept is supplied as tablets.

1.3 ROUTE OF ADMINISTRATION

- Oral

1.4 STERILE/RADIOACTIVE STATEMENT

Not applicable

1.5 QUALITATIVE AND QUANTITATIVE COMPOSITION

Active ingredient: mycophenolate mofetil.

Oral administration:

Each tablet contains 500 mg mycophenolate mofetil.

Excipients:

As registered locally.

2 CLINICAL PARTICULARS

2.1 THERAPEUTIC INDICATION(S)

CellCept in combination with corticosteroids and either cyclosporine or tacrolimus is indicated for:

- prophylaxis of acute organ rejection in adult and pediatric patients and treatment of first or refractory organ rejection in adult patients receiving allogeneic renal transplants
- prophylaxis of acute organ rejection in adult and pediatric patients receiving allogeneic cardiac transplants.
- prophylaxis of acute organ rejection in adult and pediatric patients receiving allogeneic hepatic transplants

CellCept is indicated for induction and maintenance therapy of adult and pediatric patients with Class III-V lupus nephritis (diagnosed according to International Society of Nephrology/Renal Pathology Society classification).

2.2 DOSAGE AND ADMINISTRATION

Please refer to full prescribing information for corticosteroids and either cyclosporine or tacrolimus, which are used in combination with CellCept.

Transplant patients

Standard dosage for prophylaxis of renal rejection

Adults: A dose of 1 g administered orally twice a day (daily dose of 2 g) is recommended for use in renal transplant patients. Although a dose of 1.5 g administered twice daily (daily dose of 3 g) was used in clinical trials and was shown to be safe and effective, no efficacy advantage could be established for renal transplant patients. Patients receiving 2 g/day of CellCept demonstrated an overall better safety profile compared to patients receiving 3 g/day of CellCept.

Children (aged 3 months to 18 years): The recommended dose for patients with a body surface area > 1.5 m² may be prescribed CellCept tablets at a dose of 1 g twice daily (2 g daily dose).

Standard dosage for prophylaxis of cardiac rejection

Adults: A dose of 1.5 g administered orally twice a day (daily dose of 3 g) is recommended for use in cardiac transplant patients.

Children(aged 3 months to 18 years): The recommended starting dose of CellCept powder for oral suspension is 600 mg/m² administered twice daily. If well tolerated, the dose can be increased to a maintenance dose of 900 mg/m² twice daily (maximum total daily dose of 3 g or 15 mL of the oral suspension). The dose may be individualized based on clinical assessment [200, 201]. Patients with a body surface area of 1.25 to 1.5 m² may be prescribed CellCept capsules at a dose of 750 mg twice daily (1.5 g daily dose). Patients with a body surface area > 1.5 m² may be prescribed CellCept capsules or tablets at a dose of 1 g twice daily (2 g daily dose).

Standard dosage for prophylaxis of hepatic rejection

Adults:A dose of 1 g administered intravenously (over NO LESS THAN 2 HOURS) twice a day (daily dose of 2 g) or A dose of 1.5 g orally twice a day (daily dose of 3 g) is recommended for use in hepatic transplant patients.

Children(aged 3 months to 18 years): The recommended starting dose of CellCept powder for oral suspension is 600 mg/m² administered twice daily. If well tolerated, the dose can be increased to a maintenance dose of 900 mg/m² twice daily (maximum total daily dose of 3 g or 15 mL of the oral suspension). The dose may be individualized based on clinical assessment. Patients with a body surface area of 1.25 to 1.5 m² may be prescribed CellCept capsules at a dose of 750 mg twice daily (1.5 g daily dose). Patients with a body surface area > 1.5 m² may be prescribed CellCept capsules or tablets at a dose of 1 g twice daily (2 g daily dose).

Standard dosage for treatment of first or refractory renal rejection

Adults: A dose of 1.5 g administered orally twice a day (daily dose of 3 g) is recommended for management of first or refractory rejection.

Children: No data are available for treatment of first or refractory renal rejection in pediatric renal transplant patients.

Oral administration (see 3.2.1 Pharmacokinetic Properties, Absorption)

The initial dose of CellCept should be given as soon as possible following renal, cardiac or hepatic transplantation.

Lupus nephritis patients

Standard Dosage for Induction Therapy

Adults: A dose of 750mg – 1.5g administered orally twice a day (daily dose of up to 3g) is recommended

Children: A dose of 600 mg/m² administered orally twice a day (up to a maximum of 2g daily) is recommended

Standard Dosage for Maintenance Therapy

Adults: A dose of 500mg – 1g administered orally twice a day is recommended

Children: A dose of 300 mg/m² administered orally twice a day is recommended

CellCept should be used in combination with corticosteroids. Doses should be introduced gradually and adjusted according to clinical response. Therapeutic drug monitoring could help prevent sub-therapeutic exposure ($C_{\min} \geq 3.0$ mg/L or inter-dose AUC ≥ 35 h*mg/L).

2.2.1 SPECIAL DOSAGE INSTRUCTIONS

Pediatric use

See 2.2 Dosage and administration and 3.2.5 Pharmacokinetics in Special Populations.

Geriatric use

For transplant patients, no oral dosage adjustment is recommended (see 2.4 Warnings and Precautions).

For lupus nephritis patients, no recommendation is available.

For i.v. administration of CellCept, no recommendation is available

Renal impairment

For renal transplant patients with severe chronic renal impairment (glomerular filtration rate <25 mL/min/1.73m²) outside of the immediate post-transplant period or after treatment of acute or refractory rejection, administration of doses greater than 1g twice daily should be avoided (see 3.2 Pharmacokinetic Properties).

For post-transplant patients with delayed renal graft function, no dose adjustment is recommended but patients should be carefully monitored (see 3.2 Pharmacokinetic Properties).

For cardiac or hepatic transplant patients with severe renal impairment, no data are available.

For lupus nephritis patients with GFR<30mL/min, therapeutic drug monitoring is advised.

Hepatic impairment

For renal transplant patients with severe hepatic parenchymal disease, no dose adjustments are recommended (see 3.2 Pharmacokinetic Properties).

For cardiac transplant and lupus nephritis patients with severe hepatic parenchymal disease, no data are available.

Patients with neutropenia

For patients that develop neutropenia (absolute neutrophil count $<1.3 \times 10^3/\mu\text{L}$), dosing with CellCept should be interrupted or the dose should be reduced (see 2.4 Warnings and Precautions).

2.3 CONTRAINDICATIONS

Allergic reactions to CellCept have been observed. Therefore, CellCept is contraindicated in patients with a known hypersensitivity to mycophenolate mofetil or mycophenolic acid (MPA).

CellCept is contraindicated during pregnancy due to its mutagenic and teratogenic potential (see 2.5.2 Pregnancy).

CellCept is contraindicated in women of childbearing potential not using highly effective contraceptive methods (see 2.5.1 Females and Males of Reproductive Potential).

CellCept is contraindicated in women who are breastfeeding (see 2.5.3 Lactation).

2.4 WARNINGS AND PRECAUTIONS

2.4.1 GENERAL

Neoplasms

As in all patients receiving immunosuppressive regimens involving combinations of drugs, patients receiving CellCept as part of an immunosuppressive regimen are at increased risk of developing lymphomas and other malignancies, particularly of the skin (see 2.6 Undesirable Effects). The risk appears to be related to the intensity and duration of immunosuppression rather than to the use of any specific agent].

As with all patients at an increased risk for skin cancer, exposure to sunlight and UV light should be limited by wearing protective clothing and using a sunscreen with a high protection factor.

Infections

Over suppression of the immune system can also increase susceptibility to infection including opportunistic infections, fatal infections and sepsis (see 2.6 Undesirable Effects). Such infections include latent viral reactivation, such as hepatitis B or hepatitis C reactivation, or infections caused by polyomaviruses. Cases of hepatitis due to reactivation of hepatitis B or hepatitis C have been reported in carrier patients treated with immunosuppressants. Cases of Progressive Multifocal Leukoencephalopathy (PML) associated with the JC virus, sometimes fatal, have been reported in CellCept treated patients. The reported cases generally had risk factors for PML, including immunosuppressant therapies and impairment of immune function. In immunosuppressed patients, physicians should consider PML in the differential diagnosis in patients reporting neurological symptoms and consultation with a Neurologist should be considered as clinically indicated.

BK virus-associated nephropathy has been observed during the use of CellCept in patients post renal transplant. This infection can be associated with serious outcomes, sometimes leading to renal graft loss. Patient monitoring may help detect patients at risk for BK virus-associated nephropathy. Due to the cytostatic effect of CellCept on B- and T-lymphocytes, increased severity of COVID-19 may occur. Dose reduction or discontinuation of CellCept should be considered for patients who develop evidence of BK virus-associated nephropathy, or in cases of clinically significant COVID-19.

Blood and immune system

Cases of pure red cell aplasia (PRCA) have been reported in patients treated with CellCept in combination with other immunosuppressive agents. The mechanism for mycophenolate mofetil induced PRCA is unknown; the relative contribution of other immunosuppressant and their combinations in an immunosuppression regimen are also unknown. In some cases, PRCA was found to be reversible with dose reduction or cessation of CellCept therapy. In transplant patients however reduced immunosuppression may place the graft at risk.

Patients receiving CellCept should be instructed to report immediately any evidence of infection, unexpected bruising, bleeding or any other manifestation of bone marrow depression.

Patients on CellCept should have complete blood counts weekly during the first month of treatment, twice monthly for the second and third months, then monthly through the first year. In particular, patients receiving CellCept should be monitored for neutropenia. The development of neutropenia may be related to CellCept, concomitant medications, viral infection or some combination of these causes (see 2.2.1 Special Dosage Instructions). If neutropenia develops (absolute neutrophil count $<1.3 \times 10^3/\mu\text{L}$), dosing with CellCept should be interrupted or the dose should be reduced and the patient carefully observed (see 2.2.1 Special Dosage Instructions).

Blood Donation

Patients should not donate blood during therapy and for at least 6 weeks following discontinuation of CellCept.

Vaccination

Patients should be advised that during treatment with CellCept vaccinations may be less effective and the use of live attenuated vaccines should be avoided (see 2.8 Interactions with other Medicinal Products and other Forms of Interaction). Influenza vaccination may be of value. Prescribers should refer to national guidelines for influenza vaccination.

Gastro-intestinal

CellCept has been associated with an increased incidence of digestive system adverse events, including infrequent cases of gastrointestinal tract ulceration, hemorrhage, and perforation. CellCept should be administered with caution in patients with active digestive system disease.

CellCept is an inosine monophosphate dehydrogenase (IMPDH) inhibitor; therefore, it should be avoided in patients with rare hereditary deficiency of hypoxanthine-guanine phosphoribosyl-transferase (HGPRT) such as Lesch-Nyhan and Kelley-Seegmiller syndrome.

Interactions

Caution should be exercised when switching combination therapy from regimens containing immunosuppressants, which interfere with MPA enterohepatic recirculation e.g. cyclosporineto

others devoid of this effect e.g. tacrolimus, sirolimus, belatacept, or vice versa, as this might result in changes of MPA exposure (see 2.8 Interactions with other Medicinal Products and other Forms of Interaction). Therapeutic drug monitoring of MPA may be appropriate when switching combination therapy (e.g. from cyclosporine to tacrolimus or vice versa) or to ensure adequate immunosuppression in patients with high immunological risk (e.g. risk of rejection, treatment with antibiotics, addition or removal of an interacting medication).

Drugs which interfere with MPA's enterohepatic cycle (e.g. cholestyramine, sevelamer, antibiotics) should be used with caution due to their potential to reduce the plasma levels and efficacy of CellCept (see 2.8–Interactions with other Medicinal Products and other Forms of Interaction). Sevelamer and other calcium free phosphate binders should be taken 2 hours after CellCept intake to minimise the impact on the absorption of MPA.

It is recommended that CellCept should not be administered concomitantly with azathioprine because both have the potential to cause bone marrow suppression and such concomitant administration has not been studied.

Special Populations

Geriatric population

Geriatric patients may be at an increased risk of adverse events such as certain infections (including cytomegalovirus tissue invasive disease) and possibly gastrointestinal hemorrhage and pulmonary edema, compared with younger individuals (see 2.6 Undesirable Effects).

Pregnancy and breastfeeding

CellCept is contraindicated in pregnancy and during breastfeeding (see 2.5.2 Pregnancy and 2.5.3 Lactation).

Semen Donation

Men should not donate semen during therapy and for 90 days following discontinuation of CellCept.

Patients with phenylketonuria

CellCept oral suspension contains aspartame, a source of phenylalanine (equivalent to 2.78 mg per 5 mL of oral suspension). Therefore, care should be taken if CellCept oral suspension is administered to patients with phenylketonuria.

2.4.2 DRUG ABUSE AND DEPENDENCE

There is no data available to show that CellCept has the potential for drug abuse or dependence.

2.4.3 ABILITY TO DRIVE AND USE MACHINES

CellCept may have a moderate influence on the ability to drive and use machines.

Patients should be advised to use caution when driving or using machines if they experience adverse drug reactions such as somnolence, confusion, dizziness, tremor, or hypotension during treatment with CellCept (see 2.6 Undesirable Effects).

2.5 USE IN SPECIAL POPULATIONS

2.5.1 FEMALES AND MALES OF REPRODUCTIVE POTENTIAL

Fertility

CellCept is contraindicated in women of childbearing potential not using highly effective contraceptive methods (see 2.3 Contraindications). Malformations (including anophthalmia, agnathia, and hydrocephaly) occurred in the first generation offspring of female rats treated with oral doses of mycophenolate mofetil in the absence of maternal toxicity (see 3.3.3 Impairment of Fertility). No effect was seen on the fertility of male rats treated with mycophenolate mofetil.

Pregnancy Testing

Prior to starting therapy with CellCept, female patients of childbearing potential must have a negative serum or urine pregnancy test with a sensitivity of at least 25 mIU/mL. A second test should be performed 8-10 days later. Repeat pregnancy tests should be performed during routine follow-up visits. Results of all pregnancy tests should be discussed with the patient. Patients should be instructed to consult their physician immediately should pregnancy occur.

Contraception

Females

CellCept is contraindicated in women of childbearing potential not using highly effective contraceptive methods (see 2.3 Contraindications).

Before the start of treatment, female patients of reproductive potential must be made aware of the increased risk of pregnancy loss and congenital malformations and must be counseled regarding pregnancy prevention, and planning. Women of childbearing potential should use two reliable forms of contraception simultaneously, at least one of which must be highly effective, before beginning CellCept therapy, during therapy, and for six weeks following discontinuation of therapy, unless abstinence is the chosen method of contraception.

Males

Limited clinical evidence is currently available on paternal exposure to CellCept. This evidence does not indicate an increased risk of malformations or miscarriage following paternal exposure to mycophenolate.

Non-clinical evidence shows that the dose of mycophenolate that could be transferred via the seminal fluid to a potentially pregnant partner is 30-fold lower than the concentration without teratogenic effects in animals, and 200-fold lower than the lowest teratogenic concentration in animals. Therefore, the risk of harm mediated via seminal fluid is considered negligible. However, genotoxic effects have been observed in animal studies at exposures exceeding the human therapeutic exposures by approximately 2.5-times. Thus, the risk of genotoxic effects on sperm cells cannot completely be excluded.

In absence of sufficient data to exclude a risk of harm to the fetus conceived during or directly after the treatment of the father, the following precautionary measure is recommended: sexually active male patients and/or their female partners are recommended to use effective contraception during treatment of the male patient and for at least 90 days after cessation of treatment.

2.5.2 PREGNANCY

CellCept is contraindicated during pregnancy due to its mutagenic and teratogenic potential (see 2.3 Contraindications). CellCept is a human teratogen, with an increased risk of spontaneous abortions (mainly in the first trimester) and congenital malformations in case of maternal exposure during pregnancy (see 2.6.2 Undesirable Effects, Post Marketing). In the medical literature, the risk of spontaneous abortions has been reported as 45 to 49% following mycophenolate mofetil exposure, compared to a reported rate between 12 and 33% in solid organ transplant patients treated with other immunosuppressants.

Congenital malformations (including multiple malformations in individual newborns) have been reported in 23 to 27% of live births in mycophenolate mofetil exposed pregnancies in published literature. For comparison the risk of malformations is estimated at approximately 2% of live births in the overall population and at approximately 4 to 5 % in solid organ transplant patients treated with immunosuppressants other than mycophenolate mofetil.

The following malformations were most frequently reported post-marketing, in children of patients exposed to mycophenolate mofetil in combination with other immunosuppressants during pregnancy:

- Facial malformations such as cleft lip, cleft palate, micrognathia and hypertelorism of the orbits;
- Abnormalities of the ear (e.g. abnormally formed or absent external/middle ear) and eye (e.g. coloboma, microphthalmos);
- Malformations of the fingers (e.g. polydactyly, syndactyly, brachydactyly);
- Cardiac abnormalities such as atrial and ventricular septal defects;
- Esophageal malformations (e.g. esophageal atresia);
- Nervous system malformations (such as spina bifida).

These findings were consistent with teratology studies performed in rats and rabbits where fetal resorptions and malformations occurred in absence of maternal toxicity (see 3.3.4 Reproductive Toxicity).

Labor and delivery:

The safe use of CellCept during labor and delivery has not been established.

2.5.3 LACTATION

It is not known whether the CellCept is excreted in human breast milk. Due to the potential for serious adverse reactions in nursing infants, CellCept is contraindicated during breastfeeding (see section 2.3 Contraindication).

Although the relevance to humans is unknown, studies in rats have shown mycophenolate mofetil to be excreted in milk.

2.5.4 PEDIATRIC USE

The safety and efficacy of CellCept have been established in pediatric patients 3 months and older for the prophylaxis of organ rejection of allogeneic kidney, heart or liver transplants.

Use of CellCept in pediatric kidney transplant patients is supported by evidence from adequate and well-controlled studies of CellCept in adults, with additional data from an open-label study (and follow-on study) of the safety, tolerability and efficacy of long-term oral CellCept, in 33 pediatric renal allograft recipients [39]. In addition, an open-label, pharmacokinetic and safety study of CellCept was carried out in pediatric patients after receiving allogeneic kidney transplant (100 patients, 3 months to 18 years of age).

Use of CellCept in pediatric heart transplant and liver transplant patients is supported by adequate and well-controlled studies and pharmacokinetic data in adult heart transplant and liver transplant patients. Additional supportive data include pharmacokinetic data in pediatric kidney transplant patients, pediatric liver transplant patients (9 liver transplant patients, 9 months to 12 years of age, in an open-label, pharmacokinetic and safety study) and published evidence of clinical efficacy and safety in pediatric heart transplant and pediatric liver transplant patients.

See 2.2 Dosage and Administration, 2.6 Undesirable effects, 3.1.2 Clinical / Efficacy Studies, and 3.2.5 Pharmacokinetics in Special Populations.

2.5.5 GERIATRIC USE

The safety and efficacy of CellCept have been studied in patients ranging in age from 18 to 76 years. Geriatric patients, particularly those who are receiving CellCept as part of a combination immunosuppressive regimen, may be at an increased risk of certain infections compared to other patients.

For geriatric transplant patients, no oral dosage adjustment is recommended. The safety and efficacy of administration of CellCept in lupus nephritis patients has not been studied. Likewise, the i.v. administration of CellCept in geriatric patients has not been studied for any indication.

See 2.2 Dosage and Administration, 2.4 Warnings and Precautions, 2.6 Undesirable effects, and 3.2.5 Pharmacokinetics in Special Populations.

2.5.6 RENAL IMPAIRMENT

For renal transplant patients with severe chronic renal impairment (glomerular filtration rate <25 mL/min/1.73m²) outside of the immediate post-transplant period or after treatment of acute or refractory rejection, administration of doses greater than 1g For renal transplant patients with severe hepatic parenchymal disease, no dose adjustments are recommended [3]. The safety and efficacy of CellCept in cardiac transplant and lupus nephritis patients with severe hepatic parenchymal disease, have not been studied. twice daily should be avoided.

For post-transplant patients with delayed renal graft function, no dose adjustment is recommended but patients should be carefully monitored.

The safety and efficacy of CellCept in cardiac or hepatic transplant patients with severe renal impairment, have not been studied.

For lupus nephritis patients with GFR <30 mL/min, therapeutic drug monitoring is advised.

See 2.2 Dosage and Administration and 3.2 Pharmacokinetic Properties.

2.5.7 HEPATIC IMPAIRMENT

For renal transplant patients with severe hepatic parenchymal disease, no dose adjustments are recommended. The safety and efficacy of CellCept in cardiac transplant and lupus nephritis patients with severe hepatic parenchymal disease, have not been studied.

See 2.2 Dosage and Administration and 3.2.5 Pharmacokinetics in Special Populations.

2.6 UNDESIRABLE EFFECTS

The safety profile presented in this section is based on data from both clinical trials and post marketing experience and has been shown to be consistent across transplant and lupus nephritis patient populations.

2.6.1 CLINICAL TRIALS

An estimated total of 1557 adult patients received CellCept during pivotal clinical trials in the prevention of acute organ rejection. Of these, 991 were included in the pooled renal studies ICM1866, MYC022, MYC023, 277 were included in the hepatic study MYC246, and 289 were included in the cardiac study MYC1864. Patients in all study arms also received cyclosporine and corticosteroids. Diarrhea, leukopenia, sepsis, and vomiting were among the most common and/or serious adverse drug reactions associated with the administration of CellCept in the pivotal trials. There was also evidence of a higher frequency of certain types of infection, e.g. opportunistic infections (see 2.4 Warnings and Precautions).

In the three pivotal trials for prevention of renal transplant rejection, patients receiving 2 g per day of CellCept demonstrated an overall better safety profile than patients receiving 3 g CellCept. The safety profile of CellCept in patients treated for refractory renal transplant rejection was similar to that observed in the pivotal trials for prevention of renal rejection at doses of 3 g per day. Diarrhea and leukopenia, followed by anemia, nausea, abdominal pain, sepsis, nausea and vomiting, and dyspepsia were the predominant adverse events reported more frequently in patients receiving CellCept in comparison to patients receiving i.v. corticosteroids.

The adverse event profile associated with the administration of CellCept i.v. has been shown to be similar to that observed after oral administration.

Tabulated summary of adverse drug reactions from clinical trials

Adverse drug reactions from clinical trials (Table 1) are listed by MedDRA system organ class along with their incidence. The corresponding frequency category for each adverse drug reaction is based on the following convention: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$). Due to the large differences observed in the frequency of certain ADRs across the different transplant indications, the frequency is presented separately for renal, hepatic and cardiac transplant patients.

Table 1 Summary of adverse drug reactions occurring in patients treated with CellCept in pivotal clinical trials

Adverse drug reaction (MedDRA) System Organ Class	Renal transplant n = 991		Hepatic transplant n = 277		Cardiac transplant n = 289	
	Incidence (%)	Frequency	Incidence (%)	Frequency	Incidence (%)	Frequency
Infections and infestations						
Bacterial infections	39.9	Very Common	27.4	Very Common	19.0	Very Common
Fungal infections	9.2	Common	10.1	Very Common	13.1	Very Common
Viral infections	16.3	Very Common	14.1	Very Common	31.1	Very Common
Neoplasms benign, malignant and unspecified (including cysts and polyps)						
Benign neoplasm of skin	4.4	Common	3.2	Common	8.3	Common
Neoplasm	1.6	Common	2.2	Common	4.2	Common
Skin cancer	3.2	Common	0.7	Uncommon	8.0	Common
Blood and lymphatic system disorders						
Anemia	20.0	Very Common	43.0	Very Common	45.0	Very Common
Ecchymosis	3.6	Common	8.7	Common	20.1	Very Common
Leukocytosis	7.6	Common	22.4	Very Common	42.6	Very Common
Leukopenia	28.6	Very Common	45.8	Very Common	34.3	Very Common
Pancytopenia	1.0	Common	3.2	Common	0.7	Uncommon
Pseudolymphoma	0.6	Uncommon	0.4	Uncommon	1.0	Common
Thrombocytopenia	8.6	Common	38.3	Very Common	24.2	Very Common
Metabolism and nutrition disorders						
Acidosis	3.4	Common	6.5	Common	14.9	Very Common
Hypercholesterolemia	11.0	Very Common	4.7	Common	46.0	Very Common
Hyperglycemia	9.0	Common	43.7	Very Common	48.4	Very Common
Hyperkalemia	7.3	Common	22.0	Very Common	16.3	Very Common

Hyperlipidemia	7.6	Common	8.7	Common	13.8	Very Common
Hypocalcemia	3.2	Common	30.0	Very Common	8.0	Common
Hypokalemia	7.8	Common	37.2	Very Common	32.5	Very Common
Hypomagnesemia	1.8	Common	39.0	Very Common	20.1	Very Common
Hypophosphatemia	10.8	Very Common	14.4	Very Common	8.7	Common
Weight decreased	1.0	Common	4.7	Common	6.2	Common
Psychiatric disorders						
Confusional state	1.4	Common	17.3	Very Common	14.2	Very Common
Depression	3.7	Common	17.3	Very Common	20.1	Very Common
Insomnia	8.4	Common	52.3	Very Common	43.3	Very Common
Nervous system disorders						
Dizziness	7.8	Common	16.2	Very Common	34.3	Very Common
Headache	14.8	Very Common	53.8	Very Common	58.5	Very Common
Hypertonia	3.3	Common	7.6	Common	17.3	Very Common
Paresthesia	6.3	Common	15.2	Very Common	15.6	Very Common
Somnolence	2.6	Common	7.9	Common	12.8	Very Common
Tremor	9.2	Common	33.9	Very Common	26.3	Very Common
Cardiac disorders						
Tachycardia	4.3	Common	22.0	Very Common	22.8	Very Common
Vascular disorders						
Hypertension	27.5	Very Common	62.1	Very Common	78.9	Very Common
Hypotension	4.9	Common	18.4	Very Common	34.3	Very Common
Venous thrombosis*	4.4	Common	2.5	Common	2.4	Common
Respiratory, thoracic and mediastinal disorders						
Cough	11.4	Very Common	15.9	Very Common	40.5	Very Common
Dyspnea	12.2	Very Common	31.0	Very Common	44.3	Very Common
Pleural effusion	2.2	Common	34.3	Very Common	18.0	Very Common

Gastrointestinal disorders						
Abdominal pain	22.4	Very Common	62.5	Very Common	41.9	Very Common
Colitis	1.6	Common	2.9	Common	2.8	Common
Constipation	18.0	Very Common	37.9	Very Common	43.6	Very Common
Decreased appetite	4.7	Common	25.3	Very Common	14.2	Very Common
Diarrhea	30.4	Very Common	51.3	Very Common	52.6	Very Common
Dyspepsia	13.0	Very Common	22.4	Very Common	22.1	Very Common
Esophagitis	4.9	Common	4.3	Common	9.0	Common
Flatulence	6.4	Common	18.8	Very Common	18.0	Very Common
Gastritis	4.4	Common	4.0	Common	9.3	Common
Gastrointestinal hemorrhage	2.7	Common	8.3	Common	7.6	Common
Gastrointestinal ulcer	3.1	Common	4.7	Common	3.8	Common
Ileus	2.4	Common	3.6	Common	2.4	Common
Nausea	18.4	Very Common	54.5	Very Common	56.1	Very Common
Stomatitis	1.4	Common	1.4	Common	3.5	Common
Vomiting	10.6	Very Common	32.9	Very Common	39.1	Very Common
Hepatobiliary disorders						
Blood alkaline phosphatase increased	5.2	Common	5.4	Common	9.3	Common
Blood lactate dehydrogenase increased	5.8	Common	0.7	Uncommon	23.5	Very Common
Hepatic enzyme increased	5.6	Common	24.9	Very Common	17.3	Very Common
Hepatitis	2.2	Common	13.0	Very Common	0.3	Uncommon
Skin and subcutaneous tissues disorders						
Alopecia	2.2	Common	2.2	Common	2.1	Common
Rash	6.4	Common	17.7	Very Common	26.0	Very Common
Musculoskeletal and connective tissue disorders						
Arthralgia	6.4	Common	6.1	Common	10.0	Very Common
Muscular weakness	3.0	Common	4.0	Common	13.8	Very Common
Renal and urinary disorders						

Blood creatinine increased	8.2	Common	19.9	Very Common	42.2	Very Common
Blood urea increased	0.8	Uncommon	10.1	Very Common	36.7	Very Common
Hematuria	10.0	Very Common	5.1	Common	5.2	Common
General disorders and administration site conditions						
Asthenia	10.8	Very Common	35.4	Very Common	49.1	Very Common
Chills	2.0	Common	10.8	Very Common	13.5	Very Common
Edema	21.0	Very Common	48.4	Very Common	67.5	Very Common
Hernia	4.5	Common	11.6	Very Common	12.1	Very Common
Malaise	2.4	Common	5.1	Common	9.0	Common
Pain	9.8	Common	46.6	Very Common	42.2	Very Common
Pyrexia	18.6	Very Common	52.3	Very Common	56.4	Very Common

Description of selected adverse drug reactions

Infections

All patients treated with immunosuppressants are at increased risk of bacterial, viral, and fungal infections (some of which may lead to a fatal outcome), including those caused by opportunistic agents and latent viral reactivation (see 2.4.1 Warning and Precautions). The risk increases with total immunosuppressive load (see 2.4 Warnings and Precautions). The most serious infections were sepsis and peritonitis. The most common opportunistic infections in patients receiving CellCept with other immunosuppressants were mucocutaneous candida, CMV viremia/syndrome, and herpes simplex. The proportion of patients with CMV viremia/syndrome was 13.5%.

Malignancies

Patients receiving CellCept as part of an immunosuppressive regime are at increased risk of developing lymphomas and other malignancies, particularly of the skin (see 2.4 Warnings and Precautions).

Three-year safety data in renal and cardiac transplant patients did not reveal any unexpected changes in the incidence of malignancy compared to the 1-year data. Hepatic transplant patients were followed for at least 1 year, but less than 3 years. In supportive clinical trials of treatment of refractory renal rejection, the lymphoma rate was 3.9% at an average follow-up of 42 months.

Blood and lymphatic disorders

Cytopenias, including leukopenia, anemia, thrombocytopenia and pancytopenia, are a known risk associated with mycophenolate and may lead or contribute to the occurrence of infections and hemorrhages (see 2.4 Warnings and Precautions).

Gastrointestinal

The most serious gastrointestinal disorders were ulceration and hemorrhage which are known risks associated with CellCept. Mouth, esophageal, gastric, duodenal, and intestinal ulcers often complicated by hemorrhage, as well as hematemesis, melena, and hemorrhagic forms of gastritis and colitis were commonly reported during the pivotal clinical trials. The most common gastrointestinal disorders however, were diarrhea, nausea and vomiting. Endoscopic investigation of patients with CellCept-related diarrhea have revealed isolated cases of intestinal villous atrophy (see 2.4 Warnings and Precautions).

General disorders and administration site conditions

Edema, including peripheral, face and scrotal edema, was reported very commonly during the pivotal trials. Musculoskeletal pain such as myalgia, and neck and back pain were also very commonly reported

Special Populations

Children (aged 3 months to 18 years)

The type and frequency of adverse drug reactions in a clinical study of 100 pediatric renal transplant patients aged 3 months to 18 years given 600 mg/m² mycophenolate mofetil orally twice daily, were generally similar to those observed in adult patients given 1 g CellCept twice daily. However, the following treatment-related adverse events occurred with a frequency of $\geq 10\%$ in children and were more frequent in the pediatric population, particularly in children under 6 years of age, when the frequency of treatment-related adverse events were compared to adults: diarrhea, leukopenia, sepsis, infection, and anemia.

In the open-label study in pediatric liver transplant patients and the available medical literature on pediatric liver and heart transplant patients, the type and frequency of the reported adverse reactions are consistent with those observed in pediatric patients and in adults following renal transplant.

Geriatric population (≥ 65 years)

Geriatric patients, particularly those who are receiving CellCept as part of a combination immunosuppressive regimen, may be at an increased risk of certain infections (including cytomegalovirus tissue invasive disease) and possibly gastrointestinal hemorrhage and pulmonary edema, compared to younger individuals (see 2.4 Warnings and Precautions).

2.6.2 POST-MARKETING

Adverse drug reactions in Table 2 are listed according to system organ class in MedDRA and the corresponding frequency category estimation for each adverse drug reaction is based on the following convention: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$).

Table 2 Adverse drug reactions identified from post-marketing experience

Adverse drug reaction (MedDRA) System Organ Class	Incidence (%)	Frequency category

Infections and infestations		
Protozoal infections	N/A	Uncommon ²
Neoplasms benign, malignant and unspecified (including cysts and polyps)		
Lymphoma	N/A	Uncommon ²
Lymphoproliferative disorder	N/A	Uncommon ²
Blood and lymphatic system disorders		
Aplasia pure red cell	N/A	Uncommon ²
Bone marrow failure	N/A	Uncommon ²
Gastrointestinal disorders		
Pancreatitis	1.80 ¹	Common
Immune system disorders		
Hypersensitivity	3.10 ¹	Common
Hypogammaglobulinemia	0.40 ¹	Uncommon
Respiratory, thoracic and mediastinal disorders		
Bronchiectasis	N/A	Uncommon ²
Interstitial lung disease	0.20 ¹	Uncommon
Pulmonary fibrosis	0.40 ¹	Uncommon
Vascular disorders		
Lymphocele	N/A	Uncommon ²
General disorders and administration site conditions		
De novo purine synthesis inhibitors-associated acute inflammatory syndrome	N/A	Uncommon ²

1 Highest incidence observed during the pivotal clinical trials.

2 The frequency category for ADRs observed only in the postmarketing setting is defined as the upper limit of the 95% confidence interval calculated on the basis of the total number of patients exposed to CellCept in pivotal trials.

Infections:

Serious life-threatening infections such as meningitis and infectious endocarditis have been reported occasionally and there is evidence of a higher frequency of certain types of infections such as tuberculosis and atypical mycobacterial infection.

Progressive Multifocal Leukoencephalopathy (PML) and BK virus-associated nephropathy have been reported in CellCept treated patients (see 2.4 Warnings and Precautions).

Congenital disorders and Pregnancy, puerperium, and perinatal conditions:

See 2.5.2 Pregnancy for further information.

General disorders and administration site conditions:

De novo purine synthesis inhibitors-associated acute inflammatory syndrome is a newly described paradoxical pro-inflammatory reaction associated with mycophenolate and other purine synthesis inhibitors, characterized by fever, arthralgias, arthritis, muscle pain and elevated inflammatory markers. Anecdotal literature reports showed rapid improvements following discontinuation of the drug.

2.7 OVERDOSE

Reports of overdoses with mycophenolate mofetil have been received from clinical trials and during post-marketing experience. In many of these cases no adverse events were reported. In those overdose cases in which adverse events were reported, the events fall within the known safety profile of the drug.

It is expected that an overdose of mycophenolate mofetil could possibly result in oversuppression of the immune system and increase susceptibility to infections and bone marrow suppression (see 2.4 Warnings and Precautions). If neutropenia develops, dosing with CellCept should be interrupted or the dose reduced (see 2.4 Warnings and Precautions).

MPA cannot be removed by hemodialysis. However, at high MPAG plasma concentrations (>100 µg/mL), small amounts of MPAG are removed. Bile acid sequestrants, such as cholestyramine, can remove MPA by increasing excretion of the drug (see 3.2 Pharmacokinetic Properties).

2.8 INTERACTIONS WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION

DNA Polymerase Inhibitors (acyclovir, ganciclovir)

Acyclovir: Higher MPAG (the phenolic glucuronide of MPA) and acyclovir plasma concentrations were observed when mycophenolate mofetil was administered with acyclovir than when the drugs were administered alone. Because MPAG plasma concentrations are increased in the presence of renal impairment, as are acyclovir concentrations, the potential exists for mycophenolate and acyclovir or its prodrugs e.g. valacyclovir to compete for tubular secretion, further increasing the concentrations of both drugs.

Ganciclovir: Based on the results of a single dose administration study of recommended doses of oral mycophenolate and i.v. ganciclovir and the known effects of renal impairment on the pharmacokinetics of MMF (see 3.2 Pharmacokinetic Properties and 2.4 Warnings and Precautions) and ganciclovir, it is anticipated that coadministration of these agents (which compete for mechanisms of renal tubular secretion) will result in increases in MPAG and ganciclovir concentration. No substantial alteration of MPA pharmacokinetics is anticipated and MMF dose adjustment is not required. In patients with renal impairment in whom MMF and ganciclovir or its prodrugs valganciclovir are co-administered, patients should be monitored carefully.

Antacids and proton pump inhibitors (PPIs)

Decreased mycophenolic acid (MPA) exposure has been observed when antacids, such as magnesium and aluminium hydroxides, and PPIs, including lansoprazole and pantoprazole were administered with CellCept. When comparing rates of transplant rejection or rates of graft loss between CellCept patients taking PPIs vs. CellCept patients not taking PPIs, no significant differences were seen. These data support extrapolation of this finding to all antacids because the reduction in exposure when CellCept was co-administered with magnesium and aluminium hydroxides is considerably lower than when CellCept was co-administered with PPIs.

Sequestrants

Cholestyramine: Following single-dose administration of 1.5 g of mycophenolate mofetil to normal healthy subjects pretreated with 4 g three times daily of cholestyramine for 4 days, there was a 40% reduction in the AUC of MPA. Caution should be used during concomitant administration of drugs that interfere with enterohepatic circulation (see 2.4 Warnings and Precautions).

Sevelamer: Concomitant administration of sevelamer and CellCept in adults and pediatric patients decreased the MPA C_{max} and AUC_{0-12} by 30% and 25%, respectively (see 2.4 Warnings and Precautions, Interactions).

Immunosuppressants

Cyclosporine A: CyclosporineA (CsA) pharmacokinetics were unaffected by mycophenolate mofetil. However, CsA interferes with MPA enterohepatic recycling, resulting in reduced MPA exposures by 30-50% in renal transplant patients treated with CellCept and CsA compared with patients receiving sirolimus or belatacept and similar doses of Cellcept. Conversely, changes of MPA exposure should be expected when switching patients from CsA to one of the immunosuppressants which do not interfere with MPA's enterohepatic cycle (see 2.4 Warning and Precautions).

Tacrolimus: Exposure to tacrolimus concomitantly administered with CellCept had no effect on the AUC or C_{max} of MPA in liver transplant recipients. A similar finding was observed in a recent study in kidney transplant recipients.

In renal transplant patients it was shown that the tacrolimus concentration did not appear to be altered by CellCept.

However, in hepatic transplant patients, there was an increase of approximately 20% in tacrolimus AUC when multiple doses of CellCept (1.5 g twice daily) were administered to patients taking tacrolimus.

Antibiotics

Rifampicin: After correction for dose a 70% decrease in MPA exposure (AUC_{0-12}) has been observed with concomitant rifampicin administration in a single heart-lung transplant patient. It is therefore recommended to monitor MPA exposure levels and to adjust CellCept doses accordingly to maintain clinical efficacy when the drugs are administered concomitantly.

Antibiotics eliminating β -glucuronidase-producing bacteria in the intestine (e.g. aminoglycoside, cephalosporin, fluoroquinolone, and penicillin classes of antibiotics) may interfere with MPAG/MPA enterohepatic recirculation thus leading to reduced systemic MPA exposure (see 2.4.1 Warning and Precautions, Interactions).

Information concerning the following antibiotics is available:

Ciprofloxacin or amoxicillin plus clavulanic acid: Reductions in pre-dose (trough) MPA concentrations of 54% have been reported in renal transplant recipients in the days immediately following commencement of oral ciprofloxacin or amoxicillin plus clavulanic acid. Effects tended to diminish with continued antibiotic use and cease after discontinuation. The change in pre-dose level may not accurately represent changes in overall MPA exposure; therefore, clinical relevance of these observations is unclear.

Norfloxacin and metronidazole: Norfloxacin in combination with metronidazole reduced the MPA AUC_{0-48} by 30% following a single dose of CellCept. No such effect on the systemic exposure of MPA with either of these antibiotics occurred when they were administered separately.

Trimethoprim/sulphamethoxazole: No effect on the systemic exposure of MPA (AUC, C_{max}) was seen with the combination trimethoprim/sulfamethoxazole.

Oral contraceptives

A study of coadministration of CellCept (1g twice daily) and combined oral contraceptives containing ethinylestradiol (0.02 - 0.04 mg) and levonorgestrel (0.05 - 0.20 mg), desogestrel (0.15 mg) or gestodene (0.05 - 0.10 mg) conducted in 18 women with psoriasis over 3 menstrual cycles showed no clinically relevant influence of CellCept on serum levels of progesterone, LH and FSH, thus indicating no influence of CellCept on the ovulation-suppressing action of the oral contraceptives. The pharmacokinetics of oral contraceptives were not affected to a clinically relevant degree by coadministration of CellCept (see 2.5.1 Female and Males of Reproductive Potential).

Other interactions

Concomitant administration of drugs inhibiting glucuronidation of MPA may increase MPA exposure (e.g., increase of MPA $AUC_{0-\infty}$ by 35% was observed with concomitant administration of isavuconazole). Caution is therefore recommended when administering these drugs concomitantly with CellCept.

Concomitant administration of telmisartan and CellCept resulted in an approximately 30% decrease of mycophenolic acid (MPA) concentrations. Telmisartan changes MPA's elimination by enhancing PPAR gamma (peroxisome proliferator-activated receptor gamma) expression which in turn results in an enhanced UGT1A9 expression and activity, enhancing glucuronidation. When comparing rates of transplant rejection, rates of graft loss or adverse event profiles between CellCept patients with and without concomitant telmisartan medication, no clinical consequences of the pharmacokinetic DDI were seen.

Coadministration of probenecid with mycophenolate mofetil in monkeys raises the plasma AUC of MPAG 3-fold. Thus, other drugs known to undergo renal tubular secretion may compete with MPAG and thereby raise plasma concentrations of MPAG or the other drug undergoing tubular secretion.

Live vaccines: live vaccines should not be given to patients with an impaired immune response. The antibody response to other vaccines may be diminished (see 2.4 Warnings and Precautions).

3 PHARMACOLOGICAL PROPERTIES AND EFFECTS

3.1 PHARMACODYNAMIC PROPERTIES

3.1.1 MECHANISM OF ACTION

Mycophenolate mofetil (MMF) is the 2-morpholinoethyl ester of mycophenolic acid (MPA). MPA is a potent, selective, uncompetitive and reversible inhibitor of inosine monophosphate dehydrogenase (IMPDH), and therefore inhibits the de novo pathway of guanosine nucleotide synthesis. The mechanism by which MPA inhibits the enzymatic activity of IMPDH appears to be related to the ability of MPA to structurally mimic both nicotinamide adenine dinucleotide cofactor and a catalytic water molecule. This prevents the oxidation of IMP to xanthose-5'-monophosphate which is the committed step in de novo guanosine nucleotide biosynthesis. Two IMPDH isoforms

have been identified, isoform type I, which is present in most known cells (including resting human lymphocytes) and isoform type II, which is strongly and predominantly expressed in activated human B- and T-lymphocytes. The type II isoform is nearly five times more sensitive to inhibition by MPA than is the type I isoform. MPA has more potent cytostatic effects on lymphocytes than on other cells because T- and B-lymphocytes are critically dependent for their proliferation on de novo synthesis of purines whereas other cell types can utilise salvage pathways.

In addition to its inhibition of IMPDH and the resulting deprivation of lymphocytes, MPA also influences cellular checkpoints responsible for metabolic programming of lymphocytes [198]. It has been shown, using human CD4+ T-cells, that MPA shifts transcriptional activities in lymphocytes from a proliferative state to catabolic processes relevant to metabolism and survival leading to an anergic state of T-cells, whereby the cells become unresponsive to their specific antigen

3.1.2 CLINICAL / EFFICACY STUDIES

Transplant studies

CellCept has been administered in combination with the following agents in clinical trials for the prevention of renal, cardiac and hepatic rejection episodes: antithymocyte globulin, OKT3, cyclosporine and corticosteroids. CellCept has also been administered in combination with cyclosporine and corticosteroids for the treatment of refractory renal rejection episodes. Prior to treatment with CellCept, patients may have also received antilymphocyte globulin, antithymocyte globulin and OKT3. CellCept has further been used in clinical trials together with daclizumab and tacrolimus.

Prevention of organ rejection

Adults

The safety and efficacy of CellCept in combination with corticosteroids and cyclosporine for the prevention of organ rejection were assessed in renal transplant patients in three randomised, double-blind, multicentre trials, in cardiac patients in one randomised double-blind, multicentre trial, and in hepatic patients in one randomised, double-blind, multicentre trial.

Children

The safety, pharmacokinetics and efficacy of CellCept in combination with corticosteroids and cyclosporine for the prevention of organ rejection in paediatric renal transplant patients were assessed in an open-label, multicentre study in 100 patients (aged 3 months to 18 years).

Renal transplant

Adults

The three studies compared two dose levels of oral CellCept (1 g twice daily and 1.5 g twice daily) with azathioprine (2 studies) or placebo (1 study) when administered in combination with cyclosporine and corticosteroids to prevent acute rejection episodes.

The primary efficacy endpoint was the proportion of patients in each treatment group who experienced treatment failure within the first 6 months after transplantation (defined as biopsy-proven acute rejection on treatment or the occurrence of death, graft loss or early termination from

the study for any reason without prior biopsy-proven rejection). CellCept was studied in the following three therapeutic regimens: (1) antithymocyte globulin induction / MMF or azathioprine/cyclosporine/corticosteroids, (2) MMF or azathioprine / cyclosporine/ corticosteroids, and (3) MMF or placebo/cyclosporine/corticosteroids.

CellCept, in combination with corticosteroids and cyclosporinereduced (statistically significant at the <0.05 level) the incidence of treatment failure within the first 6 months following transplantation. The following tables summarise the results of these studies. Patients who prematurely discontinued treatment were followed for the occurrence of death or graft loss, and the cumulative incidence of graft loss and patient death are summarised separately. Patients who prematurely discontinued treatment were not followed for the occurrence of acute rejection after termination. More patients receiving CellCept discontinued (without prior biopsy-proven rejection, death or graft loss) than discontinued in the control groups, with the highest rate in the CellCept 3 g/day group. Therefore, the acute rejection rates may be underestimates, particularly in the CellCept 3 g/day group.

**Renal Transplant Studies
Incidence of Treatment Failure
(Biopsy-proven Rejection or Early Termination for Any Reason)**

USA Study* (N=499 patients)	CellCept 2 g/day (n=167 patients)	CellCept 3 g/day (n=166 patients)	Azathioprine 1 to 2 mg/kg/day (n=166 patients)
All treatment failures	31.1%	31.3%	47.6%
Early termination without prior acute rejection**	9.6%	12.7%	6.0%
Biopsy-proven rejection episode on treatment	19.8%	17.5%	38.0%

* antithymocyte globulin induction/MMF or azathioprine/cyclosporine/corticosteroids

Europe/Canada/ Australia Study* (N=503 patients)	CellCept 2 g/day (n=173 patients)	CellCept 3 g/day (n=164 patients)	Azathioprine 100 to 150 mg/day (n=166 patients)
All treatment failures	38.2%	34.8%	50.0%
Early termination without prior acute rejection**	13.9%	15.2%	10.2%
Biopsy-proven rejection episode on treatment	19.7%	15.9%	35.5%

* MMF or azathioprine/cyclosporine/corticosteroids

Europe Study* (N=491 patients)	CellCept 2 g/day (n=165 patients)	CellCept 3 g/day (n=160 patients)	Placebo (n=166 patients)
All treatment failures	30.3%	38.8%	56.0%
Early termination without prior acute rejection**	11.5%	22.5%	7.2%
Biopsy-proven rejection episode on treatment	17.0%	13.8%	46.4%

* MMF or placebo/cyclosporine/corticosteroids

** Does not include death and graft loss as reason for early termination.

Cumulative incidence of 12-month graft loss and patient death are presented below. No advantage of CellCept with respect to graft loss and patient death was established. Numerically, patients receiving CellCept 2 g/day and 3 g/day experienced a better outcome than controls in all three studies; patients receiving CellCept 2 g/day experienced a better outcome than CellCept 3 g/day in two of the three studies. Patients in all treatment groups who terminated treatment early were found to have a poor outcome with respect to graft loss and patient death at 1 year.

Renal Transplant Studies
Cumulative Incidence of Combined Graft Loss
and Patient Death at 12 Months

Study	CellCept 2 g/day	CellCept 3 g/day	Control (Azathioprine or Placebo)
USA	8.5%	11.5%	12.2%
Europe/Canada/Australia	11.7%	11.0%	13.6%
Europe	8.5%	10.0%	11.5%

Children (aged 3 months to 18 years):

An open-label study of the safety, tolerability and efficacy of long-term oral CellCept was conducted in 33 paediatric renal allograft recipients. Each patient was stratified to 1 of 3 age groups (3 months to <6 years, 6 years to <12 years and 12 to 18 years) and received 1 of 3 dose levels (15 mg/kg, 23 mg/kg or 30 mg/kg twice daily). One to two-year outcomes were measured until the study was closed early as it was possible to make more specific dose recommendations. 26 of these patients then completed a follow-on study in which they could continue on a dosage of 23 mg/kg twice daily - to total approximately three years of CellCept treatment (combined duration of both studies). These studies established that the dose predicted to provide an MPA AUC_{0-12h} closest to the target exposure of 27.2 h·µg/mL was 600 mg/m², and that doses calculated based on estimated body surface area (BSA) reduced the coefficient of variation (CV) by about 10%. Therefore, dosing based on BSA is preferred over dosing based on body weight

An open-label, safety, pharmacokinetics and efficacy study of CellCept powder for oral suspension in combination with cyclosporine and corticosteroids for the prevention of renal allograft rejection was also performed in 100 pediatric patients (aged 3 months to 18 years) at centres in the US (9), Europe (5) and Australia (1). CellCept was dosed at 600 mg/m² twice daily (up to 1 g twice daily) in all age groups.

The primary efficacy endpoint was the proportion of patients experiencing an acute rejection episode in the first 6 months post-transplant. The rate of biopsy-proven rejection was similar across the age groups (3 months to <6 years, 6 years to <12 years, 12 years to 18 years). The overall biopsy-proven rejection rate at 6 months was comparable to adults. The combined incidence of graft loss (5%) and patient death (2%) at 12 months post-transplant was similar to that observed in adult renal transplant patients.

Cardiac transplant

A double-blind, randomised, comparative, parallel-group, multicentre study was performed in primary cardiac transplant recipients. The total number of patients enrolled was 650; 72 never received study drug and 578 received study drug. Patients received CellCept 1.5 g twice daily (n=289) or azathioprine 1.5 to 3 mg/kg/day (n=289), in combination with cyclosporine and corticosteroids as maintenance immunosuppressive therapy. The two primary efficacy endpoints were: (1) the proportion of patients who, after transplantation, had at least one endomyocardial biopsy-proven rejection with hemodynamic compromise, or were retransplanted or died, within the first 6 months, and (2) the proportion of patients who died or were transplanted during the first 12 months following transplantation. Patients who prematurely discontinued treatment were followed for the occurrence of allograft rejection for up to 6 months and for the occurrence of death for 1 year.

1. Rejection: No difference was established between CellCept and azathioprine (AZA) with respect to biopsy-proven rejection with hemodynamic compromise, as presented below.

Rejection at 6 Months

	All Patients		Treated Patients	
	AZA N = 323	CellCept N = 327	AZA N = 289	CellCept N = 289

Biopsy-proven rejection with hemodynamic compromise*	121 (38%)	120 (37%)	100 (35%)	92 (32%)
--	-----------	-----------	-----------	----------

* Hemodynamic compromise occurred if any of the following criteria were met: pulmonary capillary wedge pressure ≥ 20 mm or a 25% increase; cardiac index < 2.0 l/min/m² or a 25% decrease; ejection fraction $\leq 30\%$; pulmonary artery oxygen saturation $\leq 60\%$ or a 25% decrease; presence of new S₃ gallop; fractional shortening was $\leq 20\%$ or a 25% decrease; inotropic support required to manage the clinical condition.

- Survival: In the enrolled patients, there were no statistically significant differences between patients randomised to MMF and patients randomised to AZA for death and retransplantation. In patients who received study drug, the lower limit of the 97.5% confidence interval of the difference of death and retransplantation was 0.9 at 1 year, indicating that MMF was superior to AZA in these patients, as presented below.

Death or Retransplantation at 1 Year

	All Patients		Treated Patients	
	AZA N = 323	CellCept N = 327	AZA N = 289	CellCept N = 289
Death or Retransplantation	49 (15.2%)	42 (12.8%)	33 (11.4%)	18 (6.2%)
Weighted Treatment Difference	2.6%		5.3%	
Lower Limit of 97.5% one-sided Confidence Interval	-2.5%		+0.9%	

Hepatic transplant: A double-blind, randomised, comparative, parallel-group, multicentre study in primary hepatic transplant recipients was performed at 16 centres in the United States, 2 in Canada, 4 in Europe and 1 in Australia. The total number of patients enrolled was 565 and 564 received study drug. Patients either received CellCept 1g twice daily intravenously for up to 14 days followed by CellCept 1.5 g twice daily orally or azathioprine 1 - 2 mg/kg/day intravenously followed by azathioprine 1 - 2 mg/kg/day orally, in combination with cyclosporine and corticosteroids as maintenance immunosuppressive therapy. The two primary endpoints were: (1) the proportion of patients who experienced, in the first 6 months posttransplantation, one or more episodes of biopsy-proven and treated rejection or death/retransplantation, and (2) the proportion of patients who experienced graft loss (death/retransplantation) during the first 12 months posttransplantation. Patients who prematurely discontinued treatment were followed for the occurrence of allograft rejection and for the occurrence of graft loss (death/retransplantation) for 1 year. Results: In the primary (intent-to-treat) analyses CellCept in combination with corticosteroids and cyclosporine was superior to azathioprine for prevention of acute rejection ($p = 0.025$) and equivalent to azathioprine for survival.

Rejection at 6 Months / Death or Retransplantation at 1 Year

	AZA N = 287	CellCept N = 278
Biopsy proven, treated rejection at 6 months	137 (47.7%)	107 (38.5%)
Death or retransplantation at 1 year	42 (14.6%)	41 (14.7%)

Treatment of refractory organ rejection

A randomised, open-label comparison study of MMF 3g per day against intravenous corticosteroids was conducted in 150 renal transplant recipients with refractory, acute, cellular allograft rejection. The primary endpoint was the proportion of patients who were still alive with a functioning graft at 6 months after study entry.

Results: The incidence of graft loss in the control group was unexpectedly low and the primary analysis, based on the sequential probability ratio test showed a trend toward improved graft survival in the MMF group ($p=0.081$). A secondary analysis, using the Cochran-Mantel-Haenzel

test (not adjusted for sequential monitoring) suggested a 45% reduction in the incidence of graft loss or death at 6 months after study entry in the MMF arm ($p=0.062$).

	Graft Loss or Death at 6 Months	
	i.v. Steroids N = 73	CellCept N = 77
Graft loss or death at 6 months	19 (26.0%)	11 (14.3%)

Lupus nephritis studies

Many studies have investigated the use of CellCept for induction and maintenance therapy of lupus nephritis. Two large meta-analyses provide the most robust evidence to support the use of MMF in the management of lupus nephritis:

- The Cochrane Collaboration reviewed 50 randomised controlled trials including 2846 adults and children with biopsy-proven lupus nephritis Class III-V receiving immunosuppressive therapy. MMF with corticosteroids was found to be as effective as i.v. cyclophosphamide with corticosteroids in preventing death, inducing complete remission in proteinuria and achieving stable kidney function at six months with lower rates of toxicity. In maintenance therapy, MMF with corticosteroids was more effective than AZA with corticosteroids at preventing relapse with no difference in clinically important side effects.
- The National Kidney Foundation reviewed 53 randomised trials including 4,222 adults and children (from 10 years of age) with proliferative lupus nephritis receiving immunosuppressive therapy. Thirty-eight of the trials in the Cochrane review were also included in this meta-analysis. MMF with corticosteroids was as effective as i.v. cyclophosphamide with corticosteroids for inducing remission in lupus nephritis with improved tolerability. MMF with corticosteroids was superior to AZA with corticosteroids for maintaining disease remission.

The results of these meta-analyses are in line with the results of the largest randomised controlled study in lupus nephritis, the Aspreva Lupus Management Study (ALMS), which was a two-phase study investigating the efficacy of CellCept in induction and maintenance therapy of patients with Class III-V lupus nephritis. In the induction phase of the study, patients were randomly assigned to a target dose of 3g/day CellCept or 0.5 to 1.0 g/m²/month i.v. cyclophosphamide for 6 months. Both groups received prednisone, tapered from a maximum starting dose of 60mg/day. There was no significant difference in the number of patients who met the primary endpoint of a pre-specified decrease in urine protein/creatinine ratio and stabilization or improvement in serum creatinine, supporting the finding of the meta-analyses that CellCept is as effective as i.v. cyclophosphamide for induction therapy. In the 36-month maintenance phase of the study, the primary endpoint was the time to treatment failure measured as the time until the first event which was defined as death, end-stage renal disease, sustained doubling of the serum creatinine level, renal flare, or the need for rescue therapy. CellCept (2g/day) was superior to AZA (2mg/kg/day) in time to treatment failure (hazard ratio, 0.44; 95% confidence interval, 0.25 to 0.77; $P = 0.003$), supporting the finding of the meta-analyses that MMF is more effective than azathioprine at preventing relapse.

3.1.3 IMMUNOGENICITY

No data available

3.2 PHARMACOKINETIC PROPERTIES

The pharmacokinetics of MMF have been studied in renal, cardiac and hepatic transplant patients and in patients with lupus nephritis.

In general, the pharmacokinetic profile of MPA is similar in renal and in cardiac transplant patients. In the early transplant period, hepatic transplant patients receiving a 1.5 g oral MMF dose have similar MPA levels compared to renal transplant patients receiving 1 g oral or i.v. MMF. The key characteristics of MPA absorption and disposition observed in adult transplant recipients are also seen in pediatric patients with high similarity across different transplantation types. The concentration-response relationship (inhibition of IMPDH) has been found to be independent of age. The pharmacokinetic profile of MPA in lupus nephritis is similar to that reported in transplantation (including the high variability in exposure to active drug observed) but is complicated by more unpredictable changes in renal function in lupus nephritis patients.

3.2.1 ABSORPTION

Following oral administration, mycophenolate mofetil undergoes rapid and extensive absorption and complete presystemic metabolism to the active metabolite, MPA. The mean bioavailability of oral mycophenolate mofetil, based on MPA AUC, is 94%. Mycophenolate mofetil can be measured systemically during intravenous infusion; however, after oral administration it is below the limit of quantitation (0.4µg/mL).

Immediately post-transplant (<40 days) renal, cardiac and hepatic transplant patients had mean MPA AUCs approximately 30% lower and C_{max} approximately 40% lower compared to the late transplant period (3-6 months posttransplant). This is referred to as non-stationarity of MPA pharmacokinetics. MPA AUC values obtained following administration of 1 g twice daily intravenous CellCept at the recommended infusion rate to renal patients in the immediate post-transplant phase are comparable to those observed following oral dosing. In hepatic transplant patients, administration of 1 g twice daily intravenous CellCept followed by 1.5 g twice daily oral CellCept resulted in MPA AUC values similar to those found in renal transplant patients administered 1 g CellCept twice daily.

Food had no effect on the extent of absorption (MPA AUC) of mycophenolate mofetil administered at doses of 1.5 g twice daily to renal transplant patients. However, MPA C_{max} was decreased by 40% in the presence of food.

Equivalence of oral dosage forms

Bioequivalence of CellCept oral dosage forms have been evaluated. Two 500 mg tablets have been shown to be bioequivalent to four 250mg capsules. Likewise, 1g/5mL of CellCept constituted powder for oral suspension has been shown to be bioequivalent to four 250mg capsules (see 2.2 Dosage and Administration).

3.2.2 DISTRIBUTION

Secondary increases in plasma MPA concentrations are usually observed at approximately 6-12 hours post-dose, consistent with enterohepatic recirculation. A reduction of approximately 40% in the AUC of MPA is associated with coadministration of cholestyramine (4 g three times daily), consistent with interruption of enterohepatic recirculation.

At clinically relevant concentrations, MPA is 97% bound to plasma albumin. This value is dependent on renal function; changes in albumin binding after initiating therapy may explain the non-stationarity in the pharmacokinetics of MPA.

3.2.3 METABOLISM

MPA is conjugated primarily by glucuronyltransferase (isoform UGT1A9) to form the inactive phenolic glucuronide of MPA (MPAG). In vivo, MPAG is converted back to free MPA via enterohepatic recirculation. A minor acylglucuronide (AcMPAG) is also formed. AcMPAG is pharmacologically active and is suspected to be responsible for some of MMF's side effects (diarrhea, leucopenia).

3.2.4 ELIMINATION

Oral administration of radiolabelled mycophenolate mofetil resulted in complete recovery of the administered dose, with 93% of the dose recovered in the urine and 6% recovered in the feces. Most (about 87%) of a dose is excreted in the urine as MPAG. A negligible amount of drug (<1% of dose) is excreted as MPA in the urine.

Enterohepatic recirculation interferes with accurate determination of MPA's disposition parameters; only apparent values can be indicated. In healthy volunteers and patients with autoimmune disease approximate clearance values of 10.6 L/h and 8.27 L/h respectively and half-life values of 17 h were observed. In transplant patients mean clearance values were higher (range 11.9-34.9 L/h) and mean half-life values shorter (5-11 h) with little difference between renal, hepatic or cardiac transplant patients. In the individual patients, these elimination parameters vary based on type of co-treatment with other immunosuppressants, time post-transplantation, plasma albumin concentration and renal function. These factors explain why reduced exposure is seen when CellCept is co-administered with cyclosporine (see section 2.8) and why plasma concentrations tend to increase over time compared to what is observed immediately after transplantation (see non-stationarity in sections 3.2.1 and 3.2.2).

At clinically encountered concentrations, MPA and MPAG are not removed by hemodialysis. However, at high MPAG concentrations (>100 µg/mL), small amounts of MPAG are removed. By interfering with enterohepatic circulation of the drug, bile acid sequestrants, such as cholestyramine, reduce MPA AUC (see 2.7 Overdosage).

MPA's disposition depends on several transporters. Organic anion-transporting polypeptides (OATPs) and multidrug resistance-associated protein 2 (MRP2) are involved in MPA's disposition; OATP isoforms, MRP2 and breast cancer resistance protein (BCRP) are transporters associated with the glucuronides' biliary excretion. Multidrug resistance protein 1 (MDR1) is also able to transport MPA, but its contribution seems to be confined to the absorption process. In the kidney MPA and its metabolites potentially interact with renal organic anion transporters.

3.2.5 PHARMACOKINETICS IN SPECIAL POPULATIONS

Renal impairment

In a single-dose study (6 subjects per group), mean plasma MPA AUCs observed after oral dosing in subjects with severe chronic renal impairment (glomerular filtration rate <25 mL/min/1.73m²) were 28-75% higher than those observed in normal healthy subjects or subjects with lesser degrees of renal impairment. The mean single-dose MPAG AUC was 3- to 6-fold higher in subjects with severe renal impairment than in subjects with mild renal impairment and normal healthy subjects, consistent with the known renal elimination of MPAG.

Multiple dosing of mycophenolate mofetil in patients with severe chronic renal impairment has not been studied.

There is also a paucity of information available for lupus nephritis patients with severe renal impairment. Therapeutic drug monitoring in lupus nephritis patients with GFR <30 mL/min is advisable.

Patients with delayed renal graft function post-transplant

In patients with delayed renal graft function post-transplant, mean MPA AUC₀₋₁₂ was comparable to that seen in post-transplant patients without delayed renal graft function. There may be a transient increase in the free-fraction and concentration of plasma MPA in patients with delayed renal graft function. Dose adjustment of CellCept does not appear to be necessary (see 2.2.1 Special Dosage Instructions). Mean plasma MPAG AUC₀₋₁₂ was 2- to 3-fold higher than in post-transplant patients without delayed renal graft function.

In patients with primary non-functioning graft following renal transplantation, plasma concentrations of MPAG accumulated; accumulation of MPA, if any, was much smaller.

Hepatic impairment

Overall, the pharmacokinetics of MPA and MPAG were relatively unaffected by hepatic parenchymal disease in volunteers with alcoholic cirrhosis dosed with oral MMF. Effects of hepatic disease on these processes probably depend on the particular disease. Hepatic disease with predominantly biliary damage, such as primary biliary cirrhosis, may show a different effect.

Pediatric Population (aged ≤ 18 years):

Pharmacokinetic parameters were evaluated in 55 pediatric renal transplant patients (ranging from 1 year to 18 years of age) given 600 mg/m² mycophenolate mofetil orally twice daily (up to a maximum of 1 g twice daily). This dose achieved MPA AUC values similar to those seen in adult renal transplant patients receiving CellCept at a dose of 1 g twice daily in the early and late post-transplant period. MPA AUC values across age groups were similar in the early and late post-transplant period.

A comparison of dose-normalized (to 600 mg/m²) MPA AUC values in 12 pediatric kidney transplant patients less than 6 years of age at 9 months post-transplant with those values in 7 pediatric liver transplant patients [median age 17 months (range: 10 – 60 months)] at 6 months and beyond post-transplant revealed that, at the same dose, there were on average 23% lower AUC values in the pediatric liver patients compared to pediatric kidney patients. This is consistent with

the need for higher dosing in adult liver transplant patients compared to adult kidney transplant patients to achieve the same exposure.

In adult transplant patients administered the same dosage of CellCept, there is similar MPA exposure among kidney transplant and heart transplant patients. In line with the established similarity in MPA exposure between pediatric kidney transplant and adult kidney transplant patients at their respective approved doses, it could be shown that MPA exposure at the recommended dosage will be similar in pediatric heart transplant and adult heart transplant patients.

Pharmacokinetics have not been formally evaluated in pediatric lupus nephritis patients. However, in patients with systemic lupus erythematosus (some with renal involvement), similar typical MPA AUC₀₋₁₂ values following MMF administration were observed in children, adolescent, and adult patients.

Geriatric Population (≥ 65 years)

The pharmacokinetics of mycophenolate mofetil and its metabolites have not been found to be altered in geriatric transplant patients when compared to younger transplant patients.

3.3 NONCLINICAL SAFETY

The hematopoietic and lymphoid systems were the primary organs affected in toxicology studies conducted with mycophenolate mofetil in the rat, mouse, dog and monkey. These effects occurred at systemic exposure levels that are equivalent to or less than the clinical exposure at the recommended dose of 2 g/day for renal transplant recipients. Gastrointestinal effects were observed in the dog at systemic exposure levels equivalent to or less than the clinical exposure at the recommended doses. Gastrointestinal and renal effects consistent with dehydration were also observed in the monkey at the highest dose (systemic exposure levels equivalent to or greater than clinical exposure). The nonclinical toxicity profile of mycophenolate mofetil appears to be consistent with adverse events observed in human clinical trials which now provide safety data of more relevance to the patient population (see 2.6 Undesirable Effects).

3.3.1 CARCINOGENICITY

In experimental models, mycophenolate mofetil was not tumorigenic. The highest dose tested in the animal carcinogenicity studies resulted in approximately 2 – 3 times the systemic exposure (AUC or C_{max}) observed in renal transplant patients at the recommended clinical dose of 2 g/day and 1.3 – 2 times the systemic exposure (AUC or C_{max}) observed in cardiac transplant patients at the recommended clinical dose of 3 g/day.

3.3.2 GENOTOXICITY

Two genotoxicity assays (the mouse lymphoma/thymidine kinase assay and the mouse micronucleus aberration assay) indicated a potential of mycophenolate mofetil to cause chromosomal instability at severely cytotoxic dose levels. Other genotoxicity tests (the bacterial mutation assay, the yeast mitotic gene conversion assay or the Chinese hamster ovary cell chromosomal aberration assay) did not demonstrate mutagenicity.

3.3.3 IMPAIRMENT OF FERTILITY

Mycophenolate mofetil had no effect on fertility of male rats at oral doses up to 20 mg/kg/day. The systemic exposure at this dose represents 2 to 3 times the clinical exposure at the recommended clinical dose of 2 g/day in renal transplant patients and 1.3 – 2 times the clinical exposure at the recommended clinical dose of 3 g/day in cardiac transplant patients. In a female fertility and reproduction study conducted in rats, oral doses of 4.5 mg/kg/day caused malformations (including anophthalmia, agnathia, and hydrocephaly) in the first generation offspring in the absence of maternal toxicity. The systemic exposure at this dose was approximately 0.5 times the clinical exposure at the recommended clinical dose of 2 g/day for renal transplant patients and approximately 0.3 times the clinical exposure at the recommended clinical dose of 3 g/day for cardiac transplant patients. No effects on fertility or reproductive parameters were evident in the dams or in the subsequent generation.

3.3.4 REPRODUCTIVE TOXICITY

In teratology studies in rats and rabbits, fetal resorptions and malformations occurred in rats at 6 mg/kg/day (including anophthalmia, agnathia, and hydrocephaly) and in rabbits at 90 mg/kg/day (including cardiovascular and renal anomalies, such as ectopia cordis and ectopic kidneys, and diaphragmatic and umbilical hernia), in the absence of maternal toxicity. The systemic exposure at these levels was approximately equivalent to or less than 0.5 times the clinical exposure at the recommended clinical dose of 2 g/day for renal transplant patients and approximately 0.3 times the clinical exposure at the recommended clinical dose of 3 g/day for cardiac transplant patients.

Refer to section 2.5.2 Pregnancy.

4 PHARMACEUTICAL PARTICULARS

4.1 STORAGE

CellCept tablets: Do not store above 25°C.

Shelf life: 36 months

This medicine should not be used after the expiry date (EXP) shown on the pack.

4.2 SPECIAL INSTRUCTIONS FOR USE, HANDLING AND DISPOSAL

CellCept oral administration

Mycophenolate mofetil has demonstrated teratogenic effects (see 2.5.2 Pregnancy), therefore CellCept tablets should not be crushed.

4.3 PACKS

Tablets 500 mg

50

Medicine: keep out of reach of children

Current at November 2022

Tablets:

Made for F. Hoffmann-La Roche Ltd, Basel, Switzerland

by Delpharm Milano S.r.l., Segrate, Italy & Recipharm leganes, Madrid, 28914 Leganes, Spain